



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61R 31 /505, 31 /435, 31 /38, C07D 495/04, 333/78, 333/36, 333/66		A2	(11) International Publication Number: WO 99/21617
			(43) International Publication Date: 6 May 1999 (06.05.99)
(21) International Application Number: PCT/US98/20333 (22) International Filing Date: 30 September 1998 (30.09.98) (30) Priority Data: 08/959,758 29 October 1997 (29.10.97) US 09/156,077 17 September 1998 (17.09.98) US (71) Applicant (for all designated States except US): MEDCO RESEARCH, INC. [US/US]; Suite 200, 85 T. W. Alexander Drive, Research Triangle Park, NC 27709 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): BARALDI, Pier, Giovanni [IT/IT]; Via Tulipani, 73, I-66100 Ferrara (IT). (74) Agents: SAVAGE, Michael, G. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report.	
(54) Title: ALLOSTERIC ADENOSINE RECEPTOR MODULATORS			
<div style="display: flex; justify-content: space-around; align-items: flex-end;"> <div style="text-align: center;"> <p>(IA)</p> </div> <div style="text-align: center;"> <p>(IB)</p> </div> <div style="text-align: center;"> <p>(IC)</p> </div> </div>			
(57) Abstract			
<p>The present invention relates to compounds of formulas (IA, IB and IC), the preparation thereof, pharmaceutical formulations thereof, and their use in medicine as allosteric adenosine receptor modulators for uses including protection against hypoxia and ischemia induced injury and treatment of adenosine-sensitive cardiac arrhythmias.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Allosteric Adenosine Receptor Modulators

FIELD OF THE INVENTION

The present invention relates to certain thiophene derivatives and their use in the practice of medicine as allosteric modulators of adenosine receptors.

BACKGROUND OF THE INVENTION

This application is a continuation-in-part of U.S.S.N. 08/959,758, filed on October 29, 1997 by Pier G. Baraldi, entitled "Allosteric Adenosine Receptor Modulators."

Adenosine (Ado) is an autocoid (or local hormone) that modulates numerous functions in the cardiovascular and other organ systems. The actions of Ado are mediated by at least four subtypes of cell surface receptors called A_1 , A_{2a} , A_{2b} , and A_3 . Because of the ubiquity of adenosine receptors (AdoRs) throughout the human body, their indiscriminate activation may cause undesirable side effects. Therefore, it is desirable that drugs which are administered systemically to target these receptors have some degree of organ selectivity.

The overall function of Ado appears to be the regulation of the balance between oxygen (or energy) supply and consumption (or work). Ado increases oxygen supply by causing vasodilation and decreases oxygen consumption or work by inhibiting cellular functions, *e.g.*, slowing of heart rate. Consistent with this protective function, A_1 AdoR agonists, Ado uptake blockers and Ado deaminase inhibitors have been shown to reduce cellular damage and dysfunction during hypoxia and ischemia. This protective role of Ado and A_1 AdoR agonists has been shown in heart, brain, liver, and intestines. This and other potentially beneficial actions of Ado have led to increased interest in the development of Ado-related drugs targeted to ameliorate conditions such as myocardial ischemia and stroke.

However, the widespread expression of Ado receptors and the lack of sufficiently selective adenosine agonists have been a major impediment to the successful development of direct-acting AdoR agonists to exploit the cytoprotective properties of Ado.

5

A number of compounds known to modulate the action of neurotransmitters, hormones and peptides bind at sites distinct from, but functionally linked to, the primary recognition site of the respective receptors. This form of interaction between two different ligands at the same receptor protein, which may result in modulation in the form of enhancement or inhibition of each other's binding and function, is referred to as allosterism. Positive (enhancement) or negative (inhibition) allosterism are important mechanisms of action of various biologically active agents. Numerous allosteric interactions have been exploited. Among the most well known of these are the allosteric interactions between the GABA receptor and benzodiazepines; the atrial natriuretic factor (ANF) receptor and amiloride; the dextromethorphan binding site and ropizine; and the muscarinic receptor and gallamine. Allosteric modulation of the actions of Ado on the A_1 AdoR by several 2-amino-3-benzoylthiophenes on cultured cells, cardiac and brain preparations have been reported. The specificity of these compounds for A_1 AdoRs have also been demonstrated .

20

It would be advantageous to provide allosteric modulators of Ado as an alternative to direct-acting Ado agonists and nucleoside uptake blockers, preferably those which can selectively modulate the response to Ado in only those organs or localized areas of a given organ in which production of Ado is increased.

25

It is therefore an object of the present invention to provide allosteric modulators of Ado function.




It is a further object of the present invention to provide allosteric modulators of Ado function which provide a more selective therapeutic effect than direct-acting AdoR agonists.

5 It is still a further object of the present invention to provide methods of administering allosteric modulators of Ado function which limit the times and locations at which significant release of Ado occurs so that systemic side effects are minimized.

SUMMARY OF THE INVENTION

10 Compounds useful as potent, yet selective allosteric modulators of adenosine receptors, with activity as AdoR agonists, and, in some cases, AdoR antagonists, and methods of preparation and use thereof, are disclosed.

The compound have the following general formulas IA, IB, and IC:



wherein:

R_1 is hydrogen, alkyl, substituted alkyl, or haloacetyl;

R_2 , R_3 , and R_4 are independently hydrogen, halogen, alkyl, substituted alkyl, aryl, heteroaryl, heterocyclic, lower alkenyl, lower alkanoyl, amino, trifluoromethyl, amino alkyl, nitro, or cyano;

t is 0, 1, 2, or 3;

Z is NH , $N-C(X)-NH-aryl$, $NC(X)-NH-alk$, $N-C(X)-O-alk$, $N-C(X)-O-alkaryl$, $N-C(X)-O-aryl$, $N(alk)_2^+$ (and an associated pharmaceutically acceptable anion such as F^- , Cl^- , Br^- or I^-), $N-(Gr)_m(Am)_n(Alk)_p(Ar)_q$, or $CH-(Gr)_m(Am)_n(Alk)_p(Ar)_q$,

wherein

Gr is $-SO_2-$, $-C(O)O-$, or $-C(O)-$,

Am is $-CH(NH_2)-$, an amino acid residue, or an amino protected amino acid residue,

Alk is hydrogen, alkylene, substituted alkylene, alkenylene or substituted alkenylene,

Ar is aryl or substituted aryl, wherein the substituents include one or more alkyl or substituted alkyl groups or one or more nitro groups,

m is 0 or 1

n , p , and q are independently 0, 1, or 2,

provided that at least one of m , n , p , and q is other than 0;

X is O, S or N-alk

R₅ and R₆ are independently hydrogen, alkyl, substituted alkyl, or taken together form a lower alkenyl ring of 5 or 6 members,

5 provided that if R₂, R₃, and R₄ are hydrogen,
 then, both R₅ and R₆ may be neither hydrogen nor methyl;
 further provided that if R₂ and R₃ are hydrogen while R₄ is trifluoromethyl
 or if R₂ and R₄ are hydrogen while R₃ is chloro, then both R₅ and R₆ may not
 be methyl;

10

R₇ is hydrogen, alkyl, N(alk)₂, substituted alkyl or OH (and the resulting tautomeric form in which the OH is tautomerized to a carbonyl and the imine is tautomerized to an NH group);

15

Y is nitrogen, CH, C-CN or C-C(O)OR₈; and
 wherein R₈ is hydrogen, alkyl or substituted alkyl.

20

The compounds can be used in a method for allosterically modulating adenosine receptors in a mammal, including a human. The methods involve administering an effective amount of a compound of formula IA, IB, or IC sufficient to moderate adenosine receptors to the mammal. Positive allosterism results in several beneficial effects, including cardioprotection, neuroprotection, analgesia, and treatment of sleep disorders, irritable bowel syndrome, irritable bladder, urge incontinence, and glaucoma. Negative allosterism results in several beneficial effects, including the ability to treat Alzheimer's disease and

25

congestive heart failure.

30

The compounds can be used in a pharmaceutical formulation that includes a compound of formula IA, IB, or IC and one or more excipients. Various chemical intermediates can be used to prepare the compounds of formula IA, IB, or IC.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A and 1B are graphs showing the specific binding (percent of control) of the agonist [3 H]2-chloro- N^6 -cyclopentyladenosine ([3 H]CCPA) (Figure 1A) and the antagonist [3 H]8-cyclopentyl-1,3-dipropylxanthine ([3 H]CPX) (Figure 1B) as a function of concentration (log M) of the allosteric enhancer 2-amino-3-benzoyl-6-(3-methylbut-2-en-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (Compound/Example number 20) to membranes from Chinese hamster ovary ("CHO") cells expressing recombinant human adenosine A1 receptor ("huA₁AdoR").

Figures 2A-D are graphs showing the specific binding (percent of control) of the agonist [3 H]2-chloro- N^6 -cyclopentyladenosine ([3 H]CCPA) as a function of concentration (log M) of various allosteric enhancers. In Figure 2A, the circles represent results from Compound/Example 21, and the squares represent results from Compound/Example 22. In Figure 2B, the circles represent results from Compound/Example 20, and the squares represent results from Compound/Example 28. In Figure 2C, the circles represent results from Compound/Example 7, and the squares represent results from Compound/Example 9. In Figure 2D, the circles represent results from Compound/Example 10.

Figures 3A-D are graphs showing the specific binding (percent of control) of the agonist [3 H]2-chloro- N^6 -cyclopentyladenosine ([3 H]CCPA) as a function of concentration (log M) of various allosteric enhancers. In Figure 3A, the circles represent results from Compound/Example 13, and the squares represent results from Compound/Example 18. In Figure 3B, the circles represent results from Compound/Example 24, and the squares represent results from Compound/Example 5. In Figure 3C, the circles represent results from Compound/Example 27, and the squares represent results from Compound/Example 11. In Figure 3D, the circles represent results from Compound/Example 16 and the squares represent results from Compound/Example 14.

Figure 4A is a graph showing the concentration-dependent potentiation of the negative dromotropic effect (S-H interval prolongation) of adenosine by Compound 20 in guinea pig isolated hearts as a function of time (msec) vs. concentration (μM).

5 Figure 4B is a graph showing the effect of the A_1 AdoR antagonist CPX on the enhancement by Compound 20 of the negative dromotropic action of adenosine in guinea pig isolated hearts as a function of time (msec) vs. concentration (μM).

10 Figure 5 is a graph showing the effect of the compounds in Examples 1-4 on the cAMP content (pmol/mg protein) of CHO cells expressing human recombinant A_1 adenosine receptors. Bar 1 indicates the results with no CPA. Bar 2 indicates the results with 0.1 nM CPA. Bar 3 indicates the results of 0.1 nM CPA and 0.01 μM of the tested compound. Bar 4 indicates the results of 0.1 nM CPA and 0.1 μM of the tested compound. Bar 5 indicates the results of 0.1 nM CPA and 1.0 μM of the tested compound. Bar 6
15 indicates the results of 0.1 nM CPA and 10 μM of the tested compound.
For the CPA group, bars 1 and 2 are as described above. Bars 3-6 represent incubations with 0.3, 1, 3 and 10 nM CPA, respectively.

20 Figure 6-10 are graphs showing the same effect as in Figure 5, using the compounds in Examples 5-9, 10-14, 15-19, 20-24, and 25-29, respectively.

Figure 11 is a graph showing the effect of the compounds in Examples 1-4 on the cAMP content of CHO cells in the presence of CPA, as indicated as a percent change of cAMP in the presence of CPA versus the concentration of the tested compound (log M).
25 The data from Figure 5 was re-plotted for this Figure.

Figures 12-16 are graphs showing the same effect as in Figure 11, with the data from Figures 6-10, respectively, re-plotted for these Figures.

30

DETAILED DESCRIPTION OF THE INVENTION

5 The present application discloses compounds useful as potent, yet selective allosteric modulators of adenosine receptors, with activity as AdoR agonists, and in some cases, AdoR antagonists, and methods of preparation and use thereof.

10 The compounds can be used in a method for allosterically modulating adenosine receptors in a mammal, including a human. The methods involve administering an effective amount of a compound of formula IA, IB, or IC sufficient to moderate adenosine receptors to the mammal.

15 The compounds can be used in a pharmaceutical formulation that includes a compound of formula IA, IB, or IC and one or more excipients. Various chemical intermediates can be used to prepare the compounds of formula IA, IB, or IC.

Definitions

20 As used herein, a compound is an agonist of an adenosine A₁ receptor if it is able to fully inhibit adenylate cyclase (A₁) and is able to displace [¹²⁵I]-AB-MECA in a competitive binding assay.

25 As used herein, a compound is a partial agonist of an adenosine A₁ receptor if it is able to partially inhibit adenylate cyclase (A₁) and is able to displace [¹²⁵I]-AB-MECA in a competitive binding assay.

As used herein, a compound is an antagonist of an adenosine A₁ receptor if it is able to prevent the inhibition due to an agonist and is able to displace [¹²⁵I]-AB-MECA in a competitive binding assay.

As used herein, the term "alkyl" refers to monovalent straight, branched or cyclic alkyl groups preferably having from 1 to 20 carbon atoms, more preferably 1 to 10 carbon atoms ("lower alkyl") and most preferably 1 to 6 carbon atoms. This term is exemplified by groups such as methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *n*-hexyl, and the like. The terms "alkylene" and "lower alkylene" refer to divalent radicals of the corresponding alkane. Further, as used herein, other moieties having names derived from alkanes, such as alkoxyl, alkanoyl, alkenyl, cycloalkenyl, etc when modified by "lower," have carbon chains of ten or less carbon atoms. In those cases where the minimum number of carbons are greater than one, *e.g.*, alkenyl (minimum of two carbons) and cycloalkyl, (minimum of three carbons), it is to be understood that "lower" means at least the minimum number of carbons.

As used herein, the term "substituted alkyl" refers to an alkyl group, preferably of from 1 to 10 carbon atoms ("substituted lower alkyl"), having from 1 to 5 substituents, and preferably 1 to 3 substituents, selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino aminoacyl, aminoacyloxy, oxyacylamino, cyano, halogen, hydroxyl, keto, thioketo, carboxyl, carboxylalkyl, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, hydroxyamino, alkoxyamino, nitro, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, and mono- and di-alkylamino, mono- and di-(substituted alkyl)amino, mono- and di-arylamino, mono- and di-heteroarylamino, mono- and di-heterocyclic amino, and unsymmetric di-substituted amines having different substituents selected from the group consisting of alkyl, aryl, heteroaryl and heterocyclic. As used herein, other moieties having the prefix "substituted" are intended to include one or more of the substituents listed above.

As used herein, the term "alkoxy" refers to the group "alkyl-O-", where alkyl is as defined above. Preferred alkoxy groups include, by way of example, methoxy, ethoxy, *n*-

propoxy, *iso*-propoxy, *n*-butoxy, *tert*-butoxy, *sec*-butoxy, *n*-pentoxy, *n*-hexoxy, 1,2-dimethylbutoxy, and the like.

As used herein, the term "alkenyl" refers to alkenyl groups preferably having from
5 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkenyl unsaturation. Preferred alkenyl groups include ethenyl ($-\text{CH}=\text{CH}_2$), *n*-propenyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), *iso*-propenyl ($-\text{C}(\text{CH}_3)=\text{CH}_2$), and the like.

As used herein, the term "alkynyl" refers to alkynyl groups preferably having from
10 2 to 10 carbon atoms and more preferably 2 to 6 carbon atoms and having at least 1 and preferably from 1-2 sites of alkynyl unsaturation.

As used herein, the term "acyl" refers to the groups alkyl-C(O)-, substituted alkyl-C(O)-, cycloalkyl-C(O)-, substituted cycloalkyl-C(O)-, aryl-C(O)-, heteroaryl-C(O)- and
15 heterocyclic-C(O)- where alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl and heterocyclic are as defined herein.

As used herein, the term "acylamino" refers to the group -C(O)NRR where each R
is independently hydrogen, alkyl, substituted alkyl, aryl, heteroaryl, or heterocyclic
20 wherein alkyl, substituted alkyl, aryl, heteroaryl and heterocyclic are as defined herein.

As used herein, the term "aryl" refers to an unsaturated aromatic carbocyclic group
of from 6 to 14 carbon atoms having a single ring (e.g., phenyl) or multiple condensed
(fused) rings (e.g., naphthyl or anthryl). Preferred aryls include phenyl, naphthyl and the
25 like. Unless otherwise constrained by the definition for the aryl substituent, such aryl groups can optionally be substituted with from 1 to 5 substituents and preferably 1 to 3 substituents selected from the group consisting of acyloxy, hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido,
30 carboxyl, carboxylalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic,

heterocyclooxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, trihalomethyl. Preferred substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy.

5

As used herein, the term "cycloalkyl" refers to cyclic alkyl groups of from 3 to 12 carbon atoms having a single cyclic ring or multiple condensed rings. Such cycloalkyl groups include, by way of example, single ring structures such as cyclopropyl, cyclobutyl, cyclopentyl, cyclooctyl, and the like, or multiple ring structures such as adamantanyl, and the like.

10

As used herein, the terms "halo" or "halogen" refer to fluoro, chloro, bromo and iodo and preferably is either fluoro or chloro.

15

As used herein, the term "heteroaryl" refers to an aromatic carbocyclic group of from 1 to 15 carbon atoms and 1 to 4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur within at least one ring (if there is more than one ring).

20

Unless otherwise constrained by the definition for the heteroaryl substituent, such heteroaryl groups can be optionally substituted with from 1 to 5 substituents and preferably 1 to 3 substituents selected from the group consisting of acyloxy, hydroxy, acyl, alkyl, alkoxy, alkenyl, alkynyl, substituted alkyl, substituted alkoxy, substituted alkenyl, substituted alkynyl, amino, substituted amino, aminoacyl, acylamino, alkaryl, aryl, aryloxy, azido, carboxyl, carboxylalkyl, cyano, halo, nitro, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, aminoacyloxy, oxyacylamino, thioalkoxy, substituted thioalkoxy, thioaryloxy, thioheteroaryloxy, -SO-alkyl, -SO-substituted alkyl, -SO-aryl, -SO-heteroaryl, -SO₂-alkyl, -SO₂-substituted alkyl, -SO₂-aryl, -SO₂-heteroaryl, trihalomethyl. Preferred substituents include alkyl, alkoxy, halo, cyano, nitro, trihalomethyl, and thioalkoxy. Such heteroaryl groups can have a single ring (e.g., pyridyl or furyl) or multiple condensed rings (e.g., indolizinyl or benzothienyl).

25

30

"Heterocycle" or "heterocyclic" refers to a monovalent saturated or unsaturated group having a single ring or multiple condensed rings, from 1 to 15 carbon atoms and from 1 to 4 hetero atoms selected from the group consisting of nitrogen, sulfur or oxygen within the ring.

5

Unless otherwise constrained by the definition for the heterocyclic substituent, such heterocyclic groups can be optionally substituted with 1 to 5 substituents selected from the group consisting of alkyl, substituted alkyl, alkoxy, substituted alkoxy, aryl, aryloxy, halo, nitro, heteroaryl, thiol, thioalkoxy, substituted thioalkoxy, thioaryloxy, trihalomethyl, and the like. Such heterocyclic groups can have a single ring or multiple condensed rings.

10

As to any of the above groups that contain 1 or more substituents, it is understood, of course, that such groups do not contain any substitution or substitution patterns which are sterically impractical and/or synthetically non-feasible.

15

"Pharmaceutically acceptable salts" refers to pharmaceutically acceptable salts of a compound of Formulas IA, IB, or IC, which salts are derived from a variety of organic and inorganic counter ions well known in the art and include, by way of example only, sodium, potassium, calcium, magnesium, ammonium, tetraalkylammonium, and the like; and when the molecule contains a basic functionality, salts of organic or inorganic acids, such as hydrochloride, hydrobromide, tartrate, mesylate, acetate, maleate, oxalate and the like can be used as the pharmaceutically acceptable salt.

20

The term "protecting group" or "blocking group" refers to any group which when bound to one or more hydroxyl, amino or carboxyl groups of the compounds (including intermediates thereof such as the aminolactams, aminolactones, etc.) prevents reactions from occurring at these groups and which protecting group can be removed by conventional chemical or enzymatic steps to reestablish the hydroxyl, amino or carboxyl group. Preferred removable amino blocking groups include conventional substituents such

25

as t-butyloxycarbonyl (t-BOC), benzyloxycarbonyl (CBZ), and the like which can be removed by conventional conditions compatible with the nature of the product.

5 As used herein, a "negative dromotropic effect" is a decrease in the conduction velocity of the nerve tissue in the heart. As a consequence of this slow down of the conduction velocity, the S-H interval is prolonged.

Compound Preparation


10 As used herein the term "amino acid" means an alpha amino acid selected from those amino acids which naturally occur in proteins but without regard for specific stereochemical properties. The term "protected amino acid" means an amino acid in which the alpha amine group has been protected with a protecting group, as defined above. The terms "amino acid residue" and "amino acid moiety" are used synonymously herein.

15 Certain of the compounds are sufficiently basic, (*e.g.*, amino derivatives) or acidic (*e.g.*, carboxylic acid derivatives) to form salts. Pharmaceutically acceptable salts of the compounds of formulas IA, IB and IC are within the scope of the present invention. As will be understood by those skilled in the art, pharmaceutically acceptable salts include, but are not limited to, salts with inorganic acids such as hydrochloride, sulfate, phosphate, hydrobromide, and nitrate or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, methanesulfonate, p-toluenesulfonate, palmoate, salicylate, and stearate.


20

Compounds


25 The compound have the following general formulas IA, IB, and IC:



(IA)



(IB)



(IC)

wherein:

R₁ is hydrogen, alkyl, substituted alkyl, or haloacetyl;

5

R₂, R₃, and R₄ are independently hydrogen, halogen, alkyl, substituted alkyl, aryl, heteroaryl, heterocyclic, lower alkenyl, lower alkanoyl, amino, trifluoromethyl, amino alkyl, nitro, or cyano;

10

t is 0, 1, 2, or 3;

Z is NH, N-C(X)-NH-aryl, NC(X)-NH-alk, N-C(X)-O-alk, N-C(X)-O-alkaryl, N-C(X)-O-aryl, N(alk)₂ (*) (and an associated pharmaceutically acceptable anion such as F⁻, Cl⁻, Br⁻ or I⁻), N-(Gr)_m(Am)_n(Alk)_p(Ar)_q, or CH-(Gr)_m(Am)_n(Alk)_p(Ar)_q,

wherein

Gr is $-\text{SO}_2-$, $-\text{C}(\text{O})\text{O}-$, or $-\text{C}(\text{O})-$,

5

Am is $-\text{CH}(\text{NH}_2)-$, an amino acid residue, or an amino protected amino acid residue,

Alk is hydrogen, alkylene, substituted alkylene, alkenylene or substituted alkenylene,

10

Ar is aryl or substituted aryl, wherein the substituents include one or more alkyl or substituted alkyl groups or one or more nitro groups,

m is 0 or 1,

15

n, p, and q are independently 0, 1, or 2,

provided that at least one of m, n, p, and q is other than 0;

X is O, S or N-alk,

20

R_5 and R_6 are independently hydrogen, alkyl, substituted alkyl, or taken together form a lower alkenyl ring of 5 or 6 members,

provided that if R_2 , R_3 , and R_4 are hydrogen,

then, both R_5 and R_6 may be neither hydrogen nor methyl;

further provided that if R_2 and R_3 are hydrogen while R_4 is trifluoromethyl

25

or if R_2 and R_4 are hydrogen while R_3 is chloro, then both R_5 and R_6 may not be methyl;

30

R_7 is hydrogen, alkyl, $\text{N}(\text{alk})_2$, substituted alkyl or OH (and the resulting tautomeric form in which the OH is tautomerized to a carbonyl and the imine is tautomerized to an NH group);

Y is nitrogen, CH, C-CN or C-C(O)OR₈; and
 wherein R₈ is hydrogen, alkyl or substituted alkyl.

Particular compounds include compounds of formulas IA, IB, and IC wherein:

R₁ is hydrogen,

R₂, R₃, and R₄ are independently hydrogen, halogen, or trifluoromethyl,

t is 0, 1, 2, or 3,

Z is NH, N-(CH₂)₁₋₃ phenyl, N-(ethoxycarbonylmethyl), N-(2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl), N-(3-methylbut-2-en-1-yl), N-(4-methylphenylsulfonyl), N-(4-nitro-(2-phenyleth-1-yl)), or N-(benzyloxycarbonyl);

R₅ and R₆ are both hydrogen or both methyl, or R₅ and R₆ together form a cyclopentyl or cyclohexyl ring;

R₇ is hydrogen or methyl;

R₈ is ethyl.

Specific compounds are:

Compound /Example Number	<u>Compound Name</u>
2	(2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-phenyl]methanone,
5	(2-amino-3-thienyl)-(4-chlorophenyl)methanone,

Compound /Example Number	<u>Compound Name</u>
7	2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
8	2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
5	9 2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
10	2-amino-3-(4-chloro-benzoyl) 4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
10	11 2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(3-phenylprop-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
13	2-amino-3-[3-(fluoromethyl)-benzoyl]-6-(phenylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
14	2-amino-3-(4-chloro-benzoyl)-6-(2-phenyleth-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
15	15 2-amino-3-[3-(fluoromethyl)-benzoyl]-6-(2-phenyleth-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
16	2-amino-3-(4-chloro-benzoyl)-6-(3-phenylprop-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
20	18 2-amino-3-(4-chloro-benzoyl)-6-(ethoxycarbonylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
20	2-amino-3-benzoyl-6-(3-methylbut-2-en-yl) -4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
21	2-amino-3-(4-chloro-benzoyl)-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
25	22 2-amino-3-benzoyl-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,

Compound /Example Number	<u>Compound Name</u>
23	2-amino-3-benzoyl-6-[2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl]-4,5,6,7-tetrahydrothieno [2,3-c]pyridine,
24	2-amino-3-benzoyl-4,5,6,7-tetrahydrobenzo[b]thiophene,
5	25 4-phenyl-5,6,7,8-tetrahydro[1]Benzothieno[2,3-d]pyrimidine
26	2-methyl,3-ethoxycarbonyl-4-phenyl-5,6,7,8- tetrahydro[1]Benzothieno[2,3-b]pyridine
10	27 2-Amino-3-(4-bromobenzoyl)-cyclopenta[b]thiophene
28	2-amino-3-benzoyl-6-(4-methylphenylsulphonyl)-4,5,6,7- tetrahydrothieno[2,3-c]pyridine
29	2-amino-3-(4-chlorobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7- tetrahydrothieno(2,3-c)pyridine
15	30 2-amino-3-(4-chlorobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7- tetrahydrothieno(2,3-c)pyridine
31	2-amino-3-(4-chlorobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7- tetrahydrothieno(2,3-c)pyridine
32	2-amino-3-(4-iodobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7- tetrahydrothieno(2,3-c)pyridine
33	2-amino-3-(4-bromobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7- tetrahydrothieno(2,3-c)pyridine
34	2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine
20	35 2-amino-3-(4-bromobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7- tetrahydrothieno(2,3-c)pyridine
36	2-amino-3-(4-bromobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno (2,3-c)pyridine
37	2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine


Compound /Example Number	<u>Compound Name</u>
38	2-amino-3-(4-iodobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine
39	2-amino-3-(4-phenylbenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine
40	2-amino-3-(4-phenylbenzoyl)-6,6-bis(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridinium chloride
41	2-amino-3-(4-fluorobenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine
42	2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine
43	2-amino-3-(4-phenylbenzoyl)-6,6-bis(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridinium chloride
44	2-amino-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine; and
45	2-hydroxy-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine

Preparation of the Compounds

Those skilled in the art of organic chemistry will appreciate that reactive and fragile functional groups often must be protected prior to a particular reaction, or sequence of reactions, and then restored to their original forms after the last reaction is completed.

Usually groups are protected by converting them to a relatively stable derivative. For example, a hydroxyl group may be converted to an ether group and an amine group converted to an amide or carbamate. Methods of protecting and de-protecting, also known as "blocking" and "de-blocking," are well known and widely practiced in the art, *e.g.*, see T. Green, *Protective Groups in Organic Synthesis*, John Wiley, New York (1981) or

Protective Groups in Organic Chemistry, Ed. J.F.W. McOmie, Plenum Press, London (1973).




SCHEME 1

Compounds of formula IB may be conveniently prepared according to Scheme 1.

5 In Step 1 a compound of formula (II) is reacted with a compound of formula (III) in the presence of morpholine and molecular sulfur in a protic solvent, such as ethanol, at about 50° to about 65° C for about an hour to yield a compound of formula IA wherein R₁ is hydrogen.


10 Compounds of formula (IA) wherein R₁ is other than hydrogen may be prepared according to Step 2 by reacting a compound of formula (IB) from Step 1, wherein R₁ is hydrogen, with R₁X (wherein R₁ is other than hydrogen, and X is a leaving group). For a discussion of nucleophilic displacement reactions and leaving groups, see standard organic chemistry texts such as J. March, *Advanced Organic Chemistry*, Chapter 10, John Wiley & Sons, New York (1985). Compounds of formula (II) are commercially available or may be prepared by methods known to those of skill in the art. Compound of formula (III), benzophenone derivatives, may be prepared by methods known to those of skill in the art or conveniently according to Scheme 2.

15



SCHEME 2



In Scheme 2, a compound of formula (IV), a substituted acetophenone, is alpha brominated with molecular bromine in a protic, polar solvent, such as acetic acid to yield the corresponding alpha bromo compound of formula (V). The compound of formula (III) is produced by reacting the compound of formula (V) with a source of cyanide ions, such as sodium or potassium cyanide, in a polar solvent, such as water, ethanol, or a mixture thereof.



SCHEME 3


As shown in Scheme 3, a compound of formula (IA₁) wherein Z is NH may be prepared by hydrolyzing the CO-N urethane linkage of a compound of formula (VI) under acidic conditions, *e.g.*, hydrogen bromide in acetic acid.

5 In turn, a compound of formula (VI) may be prepared in a similar manner as the reaction of Scheme 1 by substituting a compound of formula (II) with a corresponding amount of a compound of formula (VII). It may be necessary to protect the carbonyl group



of the piperidinone moiety during the synthesis of a precursor compound, *e.g.*, by converting it to an ethylenedioxy derivative as seen in formula (VIII). The protecting or
 10 blocking group is removed after the synthesis of a compound of formula (VIII) to generate a compound of formula (VII).

Compounds of formula (IA) wherein Z is a substituted nitrogen, *i.e.*, N-(Gr)_m(Am)_n(Alk)_p(Ar)_q, may be prepared by nucleophilic displacement by reacting a
 15 compound of the formula X-(Gr)_m(Am)_n(Alk)_p(Ar)_q, wherein X is a leaving group (see *March, supra*), in a polar solvent in the presence of a weak base such as sodium or potassium carbonate or a tertiary amine.



SCHEME 4

According to Scheme 4, a compounds of formula (IC), can be prepared from the corresponding compound of formula (IB), wherein R_1 is hydrogen, by reacting with a compound of formula (IX) in a protic, polar solvent, such as ethanol, in the presence of a strong base such as sodium ethoxide to form the pyridine moiety. (If Y is nitrogen, then R_8' is H_2 , and if Y is CH, then R_8' is $-C(O)OR_8$). This reaction can conveniently be carried out by mixing the reactants, solvent and base at about $0^\circ C$ followed by heating at reflux for about 10 hours. A compound of formula (IC) wherein Y is nitrogen, *i.e.*, R_8 is H_2 , can be prepared from the corresponding compound of formula (IB) by reaction with a compound of formula $R_7-C(O)NH_2$, *e.g.*, formamide, if R_7 is hydrogen, at about $180^\circ C$ for about 5 hours.

Compounds of formula (IA) wherein Z is $N-(Gr)_m(Am)_n(Alk)_p(Ar)_q$, and Am is an amino acid or an amino acid with the amino group protected, and m, p, and q are 0, may be prepared by reacting the corresponding compound wherein Z is NH with a protected derivative of an amino acid. An example of a protected amino acid is BOC-tyrosine ("BOC-Tyr-OH") wherein "BOC" is $-C(O)OC(CH_3)_3$. Preferably the reaction is run in a polar, aprotic solvent, such as dimethylformamide. Preparation of BOC derivatives of amino acids are well known in the art of protein and peptide chemistry. If desired the BOC

moiety may be removed by standard means known in the art to restore the amino acid residue.

5 Compounds which include quaternary ammonium salts, for example, at the 6-position, can be prepared by reacting the amine at the desired position with excess alkyl halides using routine alkylation conditions. Compounds with urea linkages can be prepared by reacting the amine at the 6-position with the desired isocyanate. Urethanes can be prepared by reacting the amine at the 6-position with the desired alkyl chlorocarbonate (as shown, for example, in Morrison & Boyd, Organic Chemistry, Fourth Edition, Allyn & Bacon, Inc., Boston, 1983, page 840).

10

Methods of Using the Compounds

The compounds can be used for:

- 15 • Protection against hypoxia and/or ischemia induced injuries (e.g., stroke, infarction);
- Treatment of adenosine-sensitive cardiac arrhythmias;
- antinociception (i.e., analgesics);
- anticonvulsants;
- 20 • cardioprotection, short term (e.g., prior to percutaneous angioplasty (PTDA), angioplasty, and cardiac surgeries) and long term (prevention of myocardial infarction, especially in high risk patients, reduction of infarct damage, especially in high risk patients);
- treatment of congestive heart failure;
- neuroprotection: stroke prevention, stroke treatment, treatment of
- 25 Alzheimer's disease and treatment of epilepsy;
- pain management generally, including different forms of neuropathic pain, e.g., diabetic neuropathy, post herpetic neuralgia;
- antilipid uses: reduction of free fatty acids, triglycerides, glucose;

- adjunct therapy in diabetes, including insulin and non-insulin dependent diabetes mellitus: stimulation of insulin secretion from the pancreas, increase in tissue sensitivity to insulin;
- treatment of GI disorders such as diarrhea, irritable bowel disease, irritable
5 bowel syndrome, irritable bladder, incontinence such as urge incontinence;
- treatment of glaucoma;
- treatment of sleep disorders, such as sleep apnea;
- treatment of cardiac dysarrhythmias (paroxysmal supraventricular
10 tachycardia);
- use in combination with anesthesia for post surgical pain;
- treatment of inflammation;
- diagnostic uses, for example, to determine the presence of one or more of
the above described medical conditions, or in a screening assay to determine
the effectiveness of other compounds for binding to the A₁ Ado receptor
15 (i.e., through competitive inhibition as determined by various binding
assays); and
- Other indications for which A₁AdoR agonists are used.

20 The compounds can be administered via any medically acceptable means. Suitable means of administration include oral, rectal, topical or parenteral (including subcutaneous, intramuscular and intravenous) administration, although oral or parenteral administration are preferred.

25 The amount of the compound required to be effective as an allosteric modulator of an adenosine receptor will, of course, vary with the individual mammal being treated and is ultimately at the discretion of the medical or veterinary practitioner. The factors to be considered include the condition being treated, the route of administration, the nature of the formulation, the mammal's body weight, surface area, age and general condition, and the particular compound to be administered. However, a suitable effective dose is in the range

of about 0.1 $\mu\text{g/kg}$ to about 10 mg/kg body weight per day, preferably in the range of about 1 mg/kg to about 3 mg/kg per day.

5 The total daily dose may be given as a single dose, multiple doses, *e.g.*, two to six times per day, or by intravenous infusion for a selected duration. Dosages above or below the range cited above are within the scope of the present invention and may be administered to the individual patient if desired and necessary. For example, for a 75 kg mammal, a dose range would be about 75 mg to about 220 mg per day, and a typical dose would be about 150 mg per day. If discrete multiple doses are indicated, treatment might
10 typically be 50 mg of a compound given 3 times per day.

Formulations

The compounds described above are preferably administered in formulation including an active compound, *i.e.*, a compound of formula (IA), (IB) or (IC), together with
15 an acceptable carrier for the mode of administration. Suitable pharmaceutically acceptable carriers are known to those of skill in the art. The compositions can optionally include other therapeutically active ingredients, such as antibiotics, antivirals, healing promotion agents, anti-inflammatory agents, immunosuppressants, growth factors, anti-metabolites, cell adhesion molecules (CAMs), cytotoxic agents, antibodies, vascularizing agents, anti-
20 coagulants, and anesthetics/analgesics. The carrier must be pharmaceutically acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The formulations can include carriers suitable for oral, rectal, topical or parenteral
25 (including subcutaneous, intramuscular and intravenous) administration. Preferred carriers are those suitable for oral or parenteral administration.

Formulations suitable for parenteral administration conveniently include sterile aqueous preparation of the active compound which is preferably isotonic with the blood of
30 the recipient. Thus, such formulations may conveniently contain distilled water, 5%

dextrose in distilled water or saline. Useful formulations also include concentrated solutions or solids containing the compound of formula (I) which upon dilution with an appropriate solvent give a solution suitable for parental administration above.

5 For enteral administration, the compound can be incorporated into an inert carrier in discrete units such as capsules, cachets, tablets or lozenges, each containing a predetermined amount of the active compound; as a powder or granules; or a suspension or solution in an aqueous liquid or non-aqueous liquid, *e.g.*, a syrup, an elixir, an emulsion or a draught. Suitable carriers may be starches or sugars and include lubricants, flavorings,
10 binders, and other materials of the same nature.

 A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active compound in a free-flowing form, *e.g.*, a powder or granules, optionally
15 mixed with accessory ingredients, *e.g.*, binders, lubricants, inert diluents, surface active or dispersing agents. Molded tablets may be made by molding in a suitable machine, a mixture of the powdered active compound with any suitable carrier.

 A syrup or suspension may be made by adding the active compound to a
20 concentrated, aqueous solution of a sugar, *e.g.*, sucrose, to which may also be added any accessory ingredients. Such accessory ingredients may include flavoring, an agent to retard crystallization of the sugar or an agent to increase the solubility of any other ingredient, *e.g.*, as a polyhydric alcohol, for example, glycerol or sorbitol.

25 The compounds can also be administered locally by topical application of a solution, ointment, cream, gel, lotion or polymeric material (for example, a Pluronic™, BASF), which may be prepared by conventional methods known in the art of pharmacy. In addition to the solution, ointment, cream, gel, lotion or polymeric base and the active ingredient, such topical formulations may also contain preservatives, perfumes, and
30 additional active pharmaceutical agents.

Formulations for rectal administration may be presented as a suppository with a conventional carrier, e.g., cocoa butter or Witepsol S55 (trademark of Dynamite Nobel Chemical, Germany), for a suppository base.

5 Alternatively, the compound may be administered in liposomes or microspheres (or microparticles). Methods for preparing liposomes and microspheres for administration to a patient are well known to those of skill in the art. U.S. Patent No. 4,789,734, the contents of which are hereby incorporated by reference, describes methods for encapsulating biological materials in liposomes. Essentially, the material is dissolved in an aqueous
10 solution, the appropriate phospholipids and lipids added, along with surfactants if required, and the material dialyzed or sonicated, as necessary. A review of known methods is provided by G. Gregoriadis, Chapter 14, "Liposomes," Drug Carriers in Biology and Medicine, pp. 287-341 (Academic Press, 1979). Microspheres formed of polymers or proteins are well known to those skilled in the art, and can be tailored for passage through
15 the gastrointestinal tract directly into the blood stream. Alternatively, the compound can be incorporated and the microspheres, or composite of microspheres, implanted for slow release over a period of time ranging from days to months. See, for example, U.S. Patent Nos. 4,906,474, 4,925,673 and 3,625,214, the contents of which are hereby incorporated by reference.

20 Preferred microparticles are those prepared from biodegradable polymers, such as polyglycolide, polylactide and copolymers thereof. Those of skill in the art can readily determine an appropriate carrier system depending on various factors, including the desired rate of drug release and the desired dosage.

25 The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active compound into association with a carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly
30 and intimately bringing the active compound into association with a liquid carrier or a

finely divided solid carrier and then, if necessary, shaping the product into desired unit dosage form.

In addition to the aforementioned ingredients, the formulations may further include one or more optional accessory ingredient(s) utilized in the art of pharmaceutical formulations, e.g., diluents, buffers, flavoring agents, binders, surface active agents, thickeners, lubricants, suspending agents, preservatives (including antioxidants) and the like.

Determination of the Degree of Activity for the Compounds

The activity of the compounds can be readily determined using no more than routine experimentation using any of the following assays.

Binding assays.

The prototypical allosteric enhancer PD 81,723, (prepared in Example 4; see Bruns *et al.*, *Mole. Pharm.*, **38**:939 (1990), Cao *et al.*, *Gen Pharmac.* **26**:1545 (1995), and Amoah-Apraku *et al.*, *J. Pharm. Exper. Ther.* **266**(2):611(1993)) has both enhancing and inhibitory activity at the A₁AdoR. Therefore, the effect of a novel series of benzoylthiophene derivatives was determined on both the agonist [³H]CCPA and the antagonist [³H]CPX binding to membranes prepared from CHO cells stably expressing the human A₁ AdoR (CHO-huA₁ AdoR). The enhancing activity was estimated by the magnitude of the increase in [³H]CCPA binding whereas the inhibitory and (or antagonistic) activity was evaluated by the potency of the benzoylthiophene derivatives to compete for the specific binding of [³H]CPX. The method used for the preparation of the membranes of CHO cells expressing huA₁ AdoR, and the protocols for the radioligand binding assays are described by Kollias-Baker *et al.*, (*JPET*, **281**, 761(1997) and *Circ. Res.*, **75**, 961 (1994)).

Functional Assays.

In previous studies (Amoah-Apraku *et al.*, *J. Pharmacol Exp. Ther.*, **266**, 611 (1993) and Kollias-Baker, *supra*) the prototypical allosteric enhancer PD 81,723 was shown to selectively enhance A₁ AdoR-mediated prolongation of the stimulus to His (S-H) bundle interval (negative dromotropic effect) but did not increase the A_{2a}AdoR-mediated coronary vasodilation caused by Ado. Therefore, the effect of compound 20 on the negative dromotropic action of Ado in guinea pig isolated perfused hearts was determined. The guinea pig isolated perfused heart preparation and the methods for recording His bundle electrograms and measuring the S-H intervals have been previously reported.

Results

Radioligand Binding Assays

The effect of the benzoylthiophene derivative compound 20 on agonist and antagonist binding to CHO cells expressing the recombinant huA₁AdoR was investigated. Specifically, the effects of compound 20 on the binding of the agonist radioligand [³H]CCPA (2 nM) and the antagonist radioligand [³H]CPX (1 nM) to recombinant CHO-huA₁AdoR were determined. As shown in Figure 1A, the effect of compound 20 on the specific binding of [³H]CCPA was biphasic, at concentrations up to 7 μ M it increased but thereafter it decreased the specific binding of [³H]CCPA. In contrast, compound 20 did not enhance the binding of the antagonist radioligand [³H]CPX and at concentrations greater than 1 μ M decreased the specific binding of [³H]CPX, see Figure 1B. The values are mean \pm SEM of 4 hearts. Each data point represents mean \pm SEM specific binding with determinations from 2-3 experiments. Figures 2A-D and 3A-D are similar to Figure 1A but show the result of [³H]CCPA studies on other compounds.

Functional Studies

Consistent with the results of the radioligand binding assays, compound 20 enhanced the negative dromotropic effect (S-H interval prolongation) caused by Ado in a concentration-dependent manner (Figure 4A). In presence of 1 μ M Compound 20, 3 μ M adenosine caused 2° A-V block in 2 of 4 hearts. The values are mean \pm SEM of 4 guinea

pigs. For instance, 0.1, 0.5 and 1.0 μ M compound 20 enhanced the S-H prolongation induced by 3 μ M Ado by 32%, 77%, and 311%, respectively. At 1.0 μ M compound 20, the negative dromotropic effect of Ado was maximal, eliciting 2° A-V block in 2 of 4 hearts. In contrast, in the absence of compound 20, the same concentration of Ado (3 μ M) prolonged the S-H interval by 11 ± 3 msec. To demonstrate that the enhancement of the dromotropic effect of Ado by compound 20 was mediated by activation of A_1 AdoRs, prolongation of S-H interval caused by Ado in the presence of compound 20 was shown to be reversed by 10 μ M of the A_1 AdoR antagonist CPX (Figure 4B). The reversal of the effects of compound 20 by CPX establishes that the enhancement was mediated through the A_1 AdoR.

EXAMPLES

The following examples illustrate aspects of this invention but should not be construed as limitations. The symbols and conventions used in these examples are intended to be consistent with those used in the contemporary, international, chemical literature, for example, the *Journal of the American Chemical Society* ("*J.Am.Chem.Soc.*") and *Tetrahedron*.

Example 1. Preparation of (2-Amino-4,5-dimethyl-3-thienyl)-(phenyl)methanone:

Compound 1

A. General procedure for the preparation of phenacyl-bromides, the compounds of formula (V).

A solution of bromine (55 mmol) in acetic acid (50 mL) is added dropwise to a acetophenone (50 mmol), which is a compound of formula (IV), in glacial acetic (100ml) in half an hour, with stirring. The resulting suspension is heated at 50° C for an hour, and then poured into ice water (500 mL). The precipitated phenacyl bromide, a compound of formula (V), is filtered and washed with cold water three times, and finally crystallized from ethanol. (See Rather and Reid, *J.Am.Chem.Soc.* 41, 77 (1919)).

B. General procedure for the preparation of substituted benzoyl acetonitriles, the compounds of formula (III)

A solution of the phenacyl bromide as prepared in Step A, above, in ethanol is reacted with an aqueous solution of potassium cyanide dissolved in distilled water. The reaction is monitored by TLC control and during this time the solution changes color from yellow-orange to yellow-red. When the reaction is complete, crushed ice is added in a large amount and the solution is acidified with acetic acid. The precipitated corresponding benzoyl acetonitriles are filtered, washed with cold water and then air dried.

10 C. Preparation of (2-amino-4,5-dimethyl-3-thienyl)-(phenyl)methanone

A mixture of equimolar amounts of methylethyl ketone (0.01 mol), which is a compound of formula (II) wherein R_5 and R_6 are methyl, benzoyl acetonitrile (0.01 mol), which is a compound of formula (III) wherein R_2 , R_3 , and R_4 are hydrogen, sulfur (0.01 mol) and morpholine (0.01 mol) in ethanol (4 mL) was stirred and heated at 60° C for an hour (TLC control). After this time, the suspension was left standing overnight and the mixture was poured into water and the precipitated solid was extracted with ethyl acetate (3 x 50 mL). The organic layers were dried over magnesium sulfate and evaporated under vacuum. The crude product was chromatographed on silica gel column using mixtures of ethyl acetate and petroleum ether. (m.p. 140-141°C, 80% yield). $^1\text{H-NMR}$: (CDCl_3) : 1.53 (s, 3H), 2.13 (s, 3H); 6.44 (sb, 2H); 7.43-7.54 (m, 5H).

Examples 2 - 6: Compounds 2 - 6

The following compounds of formula (IB) were prepared using the procedure of Scheme 1 taught herein above, and in an analogous manner to Example 1 using appropriate precursor compounds of formula (II) and formula (III). If the desired compounds of formulas (II) and (III) are not commercially available, they may be prepared according to Example 1, Sections A and B.

2: Preparation of (2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-phenyl]

methanone (m.p. 155-157°C, 88% yield). ¹H-NMR: (CDCl₃) : 1.71 (s, 3H), 2.16 (s, 3H); 4.79 (sb, 2H); 6.03 (sb, 2H); 7.48 (s, 2H).

3: Preparation of (2-amino-4,5-dimethyl-3-thienyl)-(4-chloro-phenyl)methanone
5 (m.p. 128-130°C, 89% yield). ¹H NMR: (CDCl₃) : 1.54 (s, 3H), 2.13 (s, 3H); 6.47 (sb, 2H); 7.35-7.48 (m, 4H).

4: Preparation of (2-amino-4,5-dimethyl-3-thienyl)-[3-(trifluoromethyl)-
10 phenyl]methanone (m.p. 103-105°C, 78% yield); ¹H NMR: (CDCl₃) : 1.48 (s, 3H), 2.13 (s, 3H); 6.67 (sb, 2H); 7.54-7.75 (m, 4H).

5: Preparation of (2-amino-3-thienyl)-(4-chlorophenyl)methanone
15 (m.p. 178-180°C, 81% yield). ¹H NMR: (DMSO-d₆) 6.27 (d, 1 H), 6.72 (d, 1 H); 7.52-7.63 (m, 4H); 8.39 (sb, 2H).

6: Preparation of (2-amino-3-thienyl)-phenylmethanone
20 (m.p. 150-152°C, 75% yield). ¹H NMR: (CDCl₃) : 6.11 (d, 1H), 6.87 (d, 1H); 7.05 (sb, 2H); 7.3-7.7 (m, 5H).

Example 7. Preparation of 2-Amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridine: Compound 7

A. Preparation of 8-benzyloxycarbonyl 1,4-dioxo-8-azaspiro [4,5] decane

25 To a well-stirred and ice-cooled solution of 1,4-dioxo-8-azaspiro[4.5]decane (34.9 mmol, 5 g) in dichloromethane (200 mL) under an argon atmosphere, was added triethylamine (52.4 mmol, 7.3 mL) and then benzyloxycarbonyl chloride (42 mmol, 5.93 mL) dropwise. The suspension was stirred at room temperature for 24 hours and the precipitated solid was filtered. The organic solution was evaporated under vacuum to give an oily residue which was chromatographed on silica gel eluting with a mixture of ethyl
30 ether and petroleum ether to afford 8-benzyloxycarbonyl 1,4-dioxo-8- azaspiro [4,5]

decane in quantitative yield. ¹H-NMR (CDCl₃): 1.63 (m, 4H); 3.56 (m, 4H); 3.89 (s, 4H); 5.09 (s, 2H); 7.28 (s, 5H).

B. Preparation of 1-benzyloxycarbonyl piperidin-4-one

5 To a stirred solution of 8-benzyloxycarbonyl 1,4-dioxo-8-azaspiro[4,5] decane (0.037 mol, 10 g) in tetrahydrofuran (150 mL) was added a solution of hydrochloric acid 5% (20 mL) dropwise at room temperature. The solution was stirred for 18h (TLC control) and then evaporated under vacuum to small volume (20 mL) and neutralized with a saturated sodium bicarbonate solution. The aqueous solution was extracted with ethyl acetate (3 x 100 mL) and the organic layers were dried over sodium sulfate and finally
10 evaporated under vacuum to give 1-benzyloxycarbonyl piperidin-4-one which was substantially pure and which was used in the next step without any purification (92% yield). ¹H-NMR (CDCl₃): 1.63 (m, 4H); 3.56 (m, 4H); 5.09 (s, 2H); 7.28 (s, 5H).

15 C. Preparation of 2-Amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridine

A mixture of equimolar amounts of 1-benzyloxycarbonyl piperidine (0.01 mol), benzoyl acetonitrile (0.01 mol), sulfur (0.01 mol) and morpholine (0.01 mol) in ethanol (4 mL) was stirred and heated at 60°C for 1h (TLC control). After this time, the suspension
20 was left to stand overnight and the mixture was poured into water and the precipitated solid was extracted with ethyl acetate (3 x 50 mL). The organic layers were dried over magnesium sulfate and evaporated under vacuum. The crude product was chromatographed on a silica gel column using mixtures of ethyl acetate and petroleum ether. (m.p. 138-140°C, 80% yield). ¹H-NMR(CDCl₃) : 1.92 (m, 2H), 3.42 (t, 2H); 4.43 (s, 2H); 5.14 (s, 2H); 6.87 (sb, 2H); 7.35-7.46 (m, 5H).
25

Example 8. Preparation of 2-Amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 8

To a cooled and stirred suspension of protected 2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (0.01 mol) in acetic acid (2
30

mL), as prepared in Example 7, was added a solution of HBr (33%) in acetic acid (10 mL). After stirring at room temperature for 4h (TLC control), n-hexane was added and the resulting suspension was evaporated under vacuum to give a solid which was dissolved in water (10 mL) and neutralized with NaOH (5% solution). The precipitated solid was chromatographed on a silica gel column eluting with ethyl acetate and petroleum ether mixture to give 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine. (m.p. 160-162°C, 92% yield). ¹H-NMR (CDCl₃) : 1.86 (m, 1 H); 1.95 (m, 2H); 2.79 (t, 2H); 3.79 (s, 2H).

Example 9. Preparation of 2-Amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 9

The procedure of Example 7 was followed except that a corresponding amount of the 4-chloro-derivative of benzoyl acetonitrile was used in place of benzoyl acetonitrile to yield 2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine. (m.p. 60 -62°C, 88% yield). ¹H-NMR (CDCl₃) : 1.94 (m, 2H), 3.45 (t, 2H); 4.44 (s, 2H); 5.16 (s, 2H); 6.85 (sb, 2H); 7.36-7.45 (m, 4H).

Example 10. Preparation of 2-Amino-3-(4-chloro-benzoyl) 4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 10

The procedure of Example 8 was followed except that a corresponding amount of 2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-tetrahydro-thieno[2,3-c]pyridine, prepared as in Example 9, was used in place of 2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine. (m.p. 164-166°C, 90% yield). ¹H-NMR (CDCl₃) : 1.74 (m, 1 H); 1.89 (m, 2H); 2.84 (t, 2H); 3.82 (s, 2H); 6.85(sb, 2H), 7.29-7.48 (m, 5H).

Example 11. Preparation of 2-Amino-3-[3-(trifluoromethyl)-benzoyl]-6-(3-phenylprop-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 11

A mixture of equimolar amounts of 3-phenylpropylpiperidin-4-one (0.01 mol) (prepared by a procedure corresponding to that of Example 7, Steps A and B), 3-

trifluoromethyl benzoyl acetonitrile (0.01 mol), sulfur (0.01 mol) and morpholine (0.01 mol) in ethanol (4 mL) was stirred and heated at 60°C for 1 hour (TLC control). After this time, the suspension was allowed to stand overnight and the mixture was poured into water and the precipitated solid was extracted with ethyl acetate (3 x 50 mL). The organic layers were dried on magnesium sulfate and evaporated under vacuum. The crude product was chromatographed on silica gel column using mixtures of ethyl acetate and petroleum ether. (m.p. 176-178°C; 78% yield). ¹H-NMR (CDCl₃): 1.88-2.00 (m, 4H); 2.45-2.71 (m, 6H); 3.44 (s, 2H); 6.83 (sb, 2H); 7.17-7.48 (m, 9H).

Example 12. Preparation of 2-Amino-3-(4-chloro-benzoyl)-6-(phenylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 12

The same procedure as Example 11 was used except that a corresponding amount of benzylpiperidin-4-one was used in place of 3-phenylpropylpiperidin-4-one and a corresponding amount of 4-chlorobenzoyl acetonitrile was used in place of 3-trifluoromethyl benzoyl acetonitrile. (m.p. 155-157°C; 78% yield).

Example 13. Preparation of 2-Amino-3-[3-(fluoromethyl)-benzoyl]-6-(phenylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 13

The same procedure as Example 11 was used except that a corresponding amount of benzylpiperidin-4-one was used in place of 3-phenylpropylpiperidin-4-one. (m.p. 58-60°C; 88% yield). ¹H-NMR (CHCl₃): 1.78-1.87 (m, 2H); 2.48 (t, 2H); 3.42 (s, 2H); 3.63 (s, 2H); 7.01 (sb, 2H); 7.28-7.74 (m, 9H).

Example 14. Preparation of 2-Amino-3-(4-chloro-benzoyl)-6-(2-phenyleth-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 14

The same procedure as Example 11 was used except that a corresponding amount of phenylethylpiperidin-4-one was used in place of 3-phenylpropylpiperidin-4-one and a corresponding amount of 4-chlorobenzoyl acetonitrile was used in place of 3-trifluoromethyl benzoyl acetonitrile. (m.p. 148-150°C; 62% yield).

Example 15. Preparation of 2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(2-phenyleth-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 15

The same procedure as Example 11 was used except that a corresponding amount of benzylpiperidin-4-one was used in place of 3-phenylpropylpiperidin-4-one. (m.p. 137-138°C; 81% yield). ¹H-NMR (CHCl₃): 1.89 (m, 2H); 2.54 (t, 2H); 2.67-2.87 (m, 4H); 3.51 (s, 2H); 6.99 (sb, 2H); 7.17-7.33 (m, 5H); 7.53-7.74 (m, 4H).

Example 16. Preparation of 2-Amino-3-(4-chloro-benzoyl)-6-(3-phenylprop-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 16

The same procedure as Example 11 was used except that a corresponding amount of 4-chlorobenzoyl acetonitrile was used in place of 3-trifluoromethyl benzoyl acetonitrile. (m.p. 98-100°C; 65% yield).

Example 17. Preparation of 2-Amino-3-benzoyl-6-methyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 17

2-Amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (2 mmol), as prepared in Example 8, and methyl iodide (3 mmol) were dissolved in dry dimethylformamide (20 mL). Finely ground anhydrous potassium carbonate (1.9 g) and sodium iodide (0.2 g) were added to the solution and the resulting mixture was warmed to 65°C overnight under nitrogen. After this period (TLC control), the reaction mixture was cooled, diluted with water, extracted with diethyl ether (3 x 50 mL), and dried on sodium sulfate. The crude product was isolated and then purified by column chromatography eluting with ethyl acetate and petroleum ether solutions to give the desired compound. (m.p. 164-165°C; 77% yield).

Example 18. Preparation of 2-Amino-3-(4-chloro-benzoyl)-6-(ethoxycarbonylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 18

The same procedure as Example 17 was used except an equivalent amount of 2-amino-3-(4-chloro-benzoyl) 4,5,6,7-tetrahydrothieno[2,3-c]pyridine was used in place of

2-Amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (2 mmol), and an equivalent amount of ethoxycarbonylmethyl iodide was used in place of methyl iodide (3 mmol). (m.p. 105-106°C; 70% yield).

5 Example 19. Preparation of 2-Amino-3-benzoyl-6-(ethoxycarbonylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 19

The same procedure as Example 17 was used except an equivalent amount of ethoxycarbonylmethyl iodide was used in place of methyl iodide (3 mmol). (m.p. 115-117°C; 83% yield).

10

Example 20. Preparation of 2-Amino-3-benzoyl-6-(3-methylbut-2-en-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 20

The same procedure as Example 17 was used except an equivalent amount of dimethylallyl iodide was used in place of methyl iodide (3 mmol). (m.p. 76 - 78° C 90% yield). ¹H-NMR (CDCl₃): 1.63 (s, 3H); 1.73 (s, 3H); 1.94 (m, 2H); 2.44 (t, 2H); 3.06 (d, 2H); 3.42 (s, 2H); 5.26 (t, 1 H); 6.80 (sb, 2H); 7.35-7.50 (m, 5H).

15

Example 21. Preparation of 2-Amino-3-(4-chloro-benzoyl)-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 21

The same procedure as Example 17 was used except an equivalent amount of 2-amino-3-(4-chloro-benzoyl) 4,5,6,7-tetrahydrothieno[2,3-c]pyridine was used in place of 2-Amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (2 mmol), and an equivalent amount of p-nitrophenylethyl iodide was used in place of methyl iodide (3 mmol). (m.p. 150-152°C; 72% yield).

20

Example 22. Preparation of 2-Amino-3-benzoyl-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 22

The same procedure as Example 17 was used except an equivalent amount of p-nitrophenylethyl iodide was used in place of methyl iodide (3 mmol). (m.p. 89-91°C; 70% yield).

25

30

Example 23. Preparation of 2-Amino-3-benzoyl-6-[2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 23

To an ice-cooled and stirred solution of 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (0.775 mmol) in dry DMF (11 mL), was added BOC-Tyr-OH (0.08 mmol) and EDCI (0.08, 0.445 g) under argon atmosphere. After stirring overnight, the mixture was evaporated under vacuum to give a solid residue, which was dissolved in saturated sodium bicarbonate solution and was extracted with ethyl acetate (3 x 20 mL), then dried on magnesium sulfate. The organic layers were evaporated under vacuum to give a solid which was chromatographed on silica gel column eluting with ethyl acetate and petroleum ether solution to afford 2-amino-3-benzoyl-6-[2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine as a yellow solid.
(m.p. 143-145°C, 84% yield).

Example 24. Preparation of 2-Amino-3-benzoyl-4,5,6,7-tetrahydrobenzo[b]thiophene: Compound 24

The same procedure as Example 1 was used except a corresponding amount of cyclohexanone was used in place of methylethyl ketone.
(m.p. 150-152°C, 75% yield). ¹H-NMR (CDCl₃): 1.46-1.49 (m, 2H), 1.69-1.80 (m, 4H); 2.47-2.54 (m, 2H); 6.71 (sb, 2H); 7.37-7.50 (m, 5H).

Example 25. Preparation of 4-Phenyl-5,6,7,8-tetrahydro[1]Benzothieno[2,3-d]pyrimidine: Compound 25

A suspension of 2-amino-3-thienyl-phenylmethanone (5 mmol) in formamide (7mL) was heated at 180° C for 5 hours in an open vessel. The residue was diluted with dimethylformamide (5mL), treated with charcoal, and filtered over a small pad of Celite 503 (brand of filter aid). The cyclized compound was precipitated by addition of water (30 mL) to the filtrate and recrystallized from the same solvents.
(m.p. 135-137° C).

Example 26. Preparation of 2-Methyl-3-ethoxycarbonyl-4-phenyl-5,6,7,8-tetrahydro[1]Benzothieno[2,3-b]pyridine: Compound 26

To an ice-cooled and stirred solution of 2-amino-3-thienyl)-phenylmethanone in absolute ethanol (20mL), ethyl acetoacetate (0.055 mol) was added. To the mixture sodium ethylate (100 mg) was added at 0° C and the solution was refluxed for about 10 hours. After completion of the reaction (TLC control), the solution was evaporated under vacuum and the residue was taken up with water (50 mL) and the aqueous solution was extracted with ethyl acetate (3 x 100 mL). The combined organic layers were dried and evaporated under vacuum to give a yellow residue which was crystallized from tetrahydrofuran/hexane.
(m.p. 118-120°C).

Example 27. Preparation of 2-Amino-3-(4-bromobenzoyl)-cyclopenta[b]thiophene: Compound 27

The same procedure as Example 1 was used except a corresponding amount of cyclopentanone was used in place of methylethyl ketone and an equivalent amount of 4-bromo-benzoyl acetonitrile was used in place of benzoyl acetonitrile. (m.p. 205-206°C, 87% yield). (CDCl₃) :2.1-2.13 (m, 4H), 2.63-2.68 (m, 2H); 6.99 (sb, 2H); 7.34 (d,2H);7.53 (d, 2H).

Example 28. Preparation of 2-Amino-3-benzoyl-6-(4-methylphenylsulphonyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine: Compound 28

To a well-stirred and ice-cooled solution of 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine (0.78 mmol, 0.2 g) in dichloromethane (20 ml) under an argon atmosphere, was added triethylamine (0.162 ml) and then p-toluenesulfonyl chloride (0.93 mmol, 177 mg) portionwise. The suspension was stirred at room temperature for 24h and the precipitated solid was filtered. The organic solution was evaporated under vacuum to give a solid residue which was chromatographed on silica gel eluting with ethyl ether and petroleum ether mixture to afford 2-amino-3-benzoyl-6-(4-methylphenylsulphonyl)-

4,5,6,7-tetrahydrothieno[2,3-c]pyridine. (m.p.165-167°C, 85 yield). ¹H-NMR (CDCl₃): 1.95 (m, 2H); 2.43 (s, 3H); 3.07 (t, 2H); 3.51 (s, 2H); 6.76 (sb, 2H); 7.3-7.67 (m, 9H).

Example 29. Preparation of 4-Phenyl-thieno[2,3-d]pyrimidine: Compound 29

5 The same procedure as Example 25 was used except a corresponding amount of 2-amino-3-benzoyl-cyclopenta[b]thiophene was used in place of 2-amino-3-thienyl)-phenylmethanone (5 mmol). In turn, 2-amino-3-benzoyl-cyclopenta[b]thiophene can be prepared by the procedure of Example 1.

10 Example 30: Preparation of 2-amino-3-(4-chlorobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 30

 To a stirred solution of 2-amino-3-(4-chlorobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine) (1.02 mmol) in dry CH₂Cl₂ (20 mL), Et₃N (1.2 eq) and phenylisocyanate (1.2 eq) were added. The mixture was stirred at room temperature for 4 hours, then was
15 concentrated at reduced pressure and the residue purified by flash chromatography (EtOAc/light petroleum 3/7) to afford the desired compound as a solid (mp 216-217 °C).

Example 31: Preparation of 2-amino-3-(4-chlorobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 31

20 To a stirred solution of 2-amino-3-(4-chlorobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH₂Cl₂ (20 mL), Et₃N (1.2 equiv.) and 3-methyl-but-2-en-yl bromide were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 118-120 °C ¹H NMR (CDCl₃) 1.63 (s, 3H); 1.73 (s, 3H); 1.94 (m, 2H); 2.45 (m,
25 2H); 3.06 (d, 2H, J = 7); 3.41 (s, 2H); 5.26 (m, 1H); 6.77 (bs, 2H); 7.39 (d, 2H, J = 6.4); 7.42 (d, 2H, J = 6.4)

Example 32: Preparation of 2-amino-3-(4-chlorobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 32

To a stirred solution of 2-amino-3-(4-chlorobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH_2Cl_2 (20 mL), Et_3N (1.2 equiv.) and prop-2-en-yl bromide (1.2 eq.) were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 118-120 °C). ^1H NMR (CDCl_3) 1.96 (m, 2H); 2.47 (m, 2H); 3.12 (d, 2H, J = 6); 3.42 (s, 2H); 5.14-5.24 (m, 2H); 5.82-5.95 (m, 1H); 6.82 (bs, 2H); 7.35 (d, 2H, J = 8); 7.43 (d, 2H, J = 8)

Example 33: Preparation of 2-amino-3-(4-iodobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 33

To a stirred solution of 2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (1.02 mmol) in dry CH_2Cl_2 (20 mL), Et_3N (1.2 eq) and phenylisocyanate (1.2 eq) were added. The mixture was stirred at room temperature for 4 hours, then was concentrated at reduced pressure and the residue purified by flash chromatography (EtOAc/light petroleum 3/7) to afford the desired compound as a solid (mp 89-90 °C).

Example 34: Preparation of 2-amino-3-(4-bromobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 34

To a stirred solution of 2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (1.02 mmol) in dry CH_2Cl_2 (20 mL), Et_3N (1.2 eq) and phenylisocyanate (1.2 eq) were added. The mixture was stirred at room temperature for 4 hours, then was concentrated at reduced pressure and the residue purified by flash chromatography (EtOAc/light petroleum 3/7) to afford the desired compound as a solid (mp 89-90 °C).

Example 35: Preparation of 2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 35

Using the procedures in Examples 7 and 8, substituting p-bromobenzoyl acetonitrile for benzoyl acetonitrile, the title compound was prepared as a solid (mp 185-187 °C).

Example 36: Preparation of 2-amino-3-(4-bromobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 36

To a stirred solution of 2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH₂Cl₂ (20 mL), Et₃N (1.2 equiv.) and 3-methyl-but-2-en-yl bromide (1.2 eq.) were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 73-75 °C ¹H NMR (CDCl₃) 1.64 (s, 3H); 1.74 (s, 3H); 1.95 (m, 2H); 2.47 (m, 2H); 5.07 (d, 2H, J = 7); 3.41 (s, 2H); 5.27 (m, 1H); 6.91 (bs, 2H); 7.34 (d, 2H, J = 8.4); 7.52 (d, 2H, J = 8.4)

Example 37: Preparation of 2-amino-3-(4-bromobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 37

To a stirred solution of 2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH₂Cl₂ (20 mL), Et₃N (1.2 equiv.) and prop-2-en-yl bromide (1.2 eq.) were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 116-118°C ¹H NMR (CDCl₃) 1.96 (m, 2H); 2.47 (m, 2H); 3.12 (d, 2H, J = 6.6);

3.42 (s, 2H); 5.15-5.25 (m, 2H); 5.80-6.00 (m, 1H); 6.81 (bs, 2H); 7.36 (d, 2H, J = 8.2); 7.53 (d, 2H, J = 8.2)

Example 38: Preparation of 2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 38

Using the procedures in Examples 7 and 8, substituting p-iodobenzoyl acetonitrile for benzoyl acetonitrile, the title compound was prepared as a solid mp 230-231 °C

Example 39: Preparation of 2-amino-3-(4-iodobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 39

To a stirred solution of 2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH_2Cl_2 (20 mL), Et_3N (1.2 equiv.) and 3-methyl-but-2-en-yl bromide (1.2 eq.) were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 148-150 °C).

Example 40: Preparation of 2-amino-3-(4-phenylbenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 40

2-Amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine was reacted with benzyloxycarbonyl chloride and triethylamine to afford the title compound as a solid (mp 83-85°C). ^1H NMR (CDCl_3) 2.05 (m, 2H); 3.45 (m, 2H); 4.46 (s, 2H); 5.15 (s, 2H); 6.72 (bs, 2H) 7.35-7.66 (m, 14H)

Example 41: Preparation of 2-amino-3-(4-phenylbenzoyl)-6,6-bis(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridinium chloride: Compound 41

To a stirred solution of 2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH_2Cl_2 (20 mL), Et_3N (1.2 equiv.) and prop-2-en-yl bromide (2.4 eq.) were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 183-185 °C). ^1H NMR (CDCl_3) 1.82 (s, 6H); 1.85 (s, 6H); 2.30 (m, 2H); 3.70 (m, 2H); 4.07 (m, 4H); 4.75 (s, 2H); 5.26 (m, 2H); 7.40-8.50 (m, 11H).

Example 42: Preparation of 2-amino-3-(4-fluorobenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 42

Using the procedures in Examples 7 and 8, substituting p-fluorobenzoyl acetonitrile for benzoyl acetonitrile, 2-amino-3-(4-fluorobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine was prepared. This compound was reacted with benzyloxycarbonyl chloride to yield the title compound as a solid (mp 90-92°C). ^1H NMR (CDCl_3) 1.96 (m, 2H); 3.45 (m, 2H); 4.45 (s, 2H); 5.15 (s, 2H); 6.71 (bs, 2H); 7.05-7.52 (m, 9H).

Example 43: Preparation of 2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine: Compound 43

Using the procedures in Examples 7 and 8, substituting p-phenylbenzoyl acetonitrile for benzoyl acetonitrile, the title compound was prepared as a solid (mp 185-187 °C). ¹H NMR (CDCl₃) 1.75 (m, 2H); 2.40 (bs, 1H); 2.6 (m, 2H); 3.59 (s, 2H); 7.44-7.776 (m, 9H); 8.16 (bs, 2H).

Example 44: Preparation of 2-amino-3-(4-phenylbenzoyl)-6,6-bis(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridinium chloride: Compound 44

To a stirred solution of 2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine (0.7 mmol) in dry CH₂Cl₂ (20 mL), Et₃N (1.2 equiv.) and prop-2-en-yl bromide (2.4 eq.) were added. The solution was stirred at room temperature for 4 hours, then concentrated and the residue purified by crystallization to afford the title compound as a solid (mp 149-151°C).

Example 45: Preparation of 2-amino-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine: Compound 45

The title compound was prepared by reacting 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine, as prepared in Example 8, with the carbanion of malonodinitrile to yield a solid (mp 152-154 °C).

Example 46: Preparation of 2-hydroxy-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine: Compound 46

The title compound was prepared by reacting 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine, as prepared in Example 8, with the carbanion of ethylcyanoacetate to yield a solid (mp 206°C).

Example 47. Pharmaceutical Formulations

(A) Transdermal System - for 1000 patches

Ingredients	Amount
Active compound	100g
Silicone fluid	450g
Colloidal silicon dioxide	2g

5

The silicone fluid and active compound are mixed together and the colloidal silicone dioxide is added to increase viscosity. The material is then dosed into a subsequent heat sealed polymeric laminate including the following: polyester release liner, skin contact adhesive composed of silicone or acrylic polymers, a control membrane which is a polyolefin, and an impermeable backing membrane made of a polyester multilaminate. The resulting laminated sheet is then cut into 10 sq. cm patches

10

(B) Oral Tablet - For 1000 Tablets

Ingredients	Amount
Active compound	50g
Starch	50g
Magnesium Stearate	5g

15

The active compound and the starch are granulated with water and dried. Magnesium stearate is added to the dried granules and the mixture is thoroughly blended. The blended mixture is compressed into tablets.

20

(C) Injection - for 1000, 1mL Ampules

Ingredients	Amount
Active compound	10g
Buffering Agents	q.s.
Propylene glycol	400mg
Water for injection	q.s.1000mL

25

The active compound and buffering agents are dissolved in the propylene glycol at about 50°C. The water for injection is then added with stirring and the resulting solution is filtered, filled into ampules, sealed and sterilized by autoclaving.

5 (D) Continuous Injection - for 1000 mL

Ingredients	Amount
Active compound	10g
Buffering agents	q.s.
Water for injection	q.s.1000mL

10

Example 48: Evaluation of Compounds

The compounds described in Examples 1-29 above were assayed for their allosteric enhancing ability at the adenosine A₁ receptor. These are listed as T1 through T29 below.

15 Materials and Methods

Compounds T1 through T29 were provided by Medco Research, Inc. The adenosine A₁ receptor agonist N⁶ cyclopentyladenosine (CPA) and the adenylyl cyclase activator forskolin were purchased from Research Biochemicals, Inc. Rolipram was a gift from Berlex Labs. Adenosine deaminase was purchased from Sigma Chemical. Ham's F-12 culture medium and fetal bovine serum were purchased from GIBCO Life Technologies. Cell culture plasticware and antibiotic G-418 were from Fisher Scientific.

20

The preparation chosen for the assay of the compounds was the Chinese hamster ovary (CHO) cell expressing human recombinant adenosine A₁ receptors at a density of around 8000 fmol/mg protein. These cells were cultured using known techniques (Shryock et al., Mol. Pharmacol., 1998, 53:886-893, the contents of which are hereby incorporated by reference).

25

Protocol

The effect of each compounds on cAMP content of cultured CHO cells was determined in the presence of forskolin (1-1.5 μ M), rolipram (20 μ M), the adenosine receptor agonist CPA (0.05-0.1 nM) and adenosine deaminase (2 U/mL). Forskoline was used to increase the activity of adenylyl cyclase and the content of cAMP in cells, and
5 rolipram was used to inhibit the activity of cAMP phosphodiesterases that degrade cAMP. Adenosine deaminase was used to degrade endogeneous adenosine in the incubation medium.

To begin an experiment, CHO cells grown in individual wells of 12-well culture
10 plates were washed once with Hank's buffered saline solution to remove growth medium. The Hank's solution was itself then removed and replaced with fresh Hank's solution at 36° C containing forskolin, rolipram, CPA, adenosine deaminase, and the compound to be assayed. After an incubation of six minute duration, this solution was removed and replaced with 50 mM hydrochloric acid to terminate the effect of the drug, lyse the cells,
15 and prevent further enzymatic formation and metabolism of cAMP. The cAMP content of acid extracts of cells was determined by radioimmunoassay as previously described (Shryock et al., *Mol. Pharmacol.*, 1998, 53:886-893). In each experiment, 4-5 compounds were tested in parallel, at each of four concentrations, 0.01, 0.1, 1 and 10 μ M. As a control, the effect of CPA (0.1-10 nM) was determined in each experiment. Protein content of cell
20 samples was measured by the method of Bradford using a kit from Bio-Rad with albumin as a reference standard.

Results

The compounds acted to both enhance and antagonize the effect of the adenosine A₁
25 receptor agonist, CPA, on CHO cells expressing adenosine A₁ receptors. The effects of all twenty nine compounds are shown in Figures 5-10 (bar graphs) and Figures 11-16 (concentration response plots). Compounds T3, T5, T7, T9, T13, T19 and T21 decreased cAMP content in the presence of a low concentration of CPA (0.05-0.1 nM). These compounds are thus indicated by the CHO cell assay to be allosteric enhancers of the action




of an adenosine A₁ receptor agonist. Compounds T5, T7, T9, and T13 appear to be the best enhancers. Compound T7 had the highest potency and efficacy (maximal effect).

5 Several compounds (i.e., T11, T12, T23, T24, and T27) acted as antagonists of the action of CPA. Two compounds (T6 and T15) at a low concentration enhanced the action of CPA, but at a higher concentration antagonized the action of CPA.

10 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed is:

1. The use of a compound in the preparation of a medicament for providing cardioprotection, neuroprotection, pain management, reduction of free fatty acids, triglycerides, or glucose levels, adjunct therapy in diabetes, treatment of GI disorders, treatment of glaucoma; treatment of sleep disorders; treatment of cardiac disarrhythmias (peroxysmal supraventricular tachycardia, treatment of congestive heart failure or treatment of inflammation, wherein the compound is selected from a group consisting of compounds of formulas IA, IB, and IC:



10 wherein:

R₁ is hydrogen, alkyl, substituted alkyl, or haloacetyl;

R_2 , R_3 , and R_4 are independently hydrogen, halogen, alkyl, substituted alkyl, aryl, heteroaryl, heterocyclic, lower alkenyl, lower alkanoyl, amino, trifluoromethyl, amino alkyl, nitro, or cyano;

5 t is 0, 1, 2, or 3;

Z is NH , $N-C(X)-NH-aryl$, $NC(X)-NH-alk$, $N-C(X)-O-alk$, $N-C(X)-O-alkaryl$, $N-C(X)-O-aryl$, $N(alk)_2$ (*) (and an associated pharmaceutically acceptable anion such as F^- , Cl^- , Br^- or I^-), $N-(Gr)_m(Am)_n(Alk)_p(Ar)_q$, or $CH-(Gr)_m(Am)_n(Alk)_p(Ar)_q$,

10 wherein

Gr is $-SO_2-$, $-C(O)O-$, or $-C(O)-$,

Am is $-CH(NH_2)-$, an amino acid residue, or an amino protected amino acid residue,

15

Alk is hydrogen, alkylene, substituted alkylene, alkenylene or substituted alkenylene,

20

Ar is aryl or substituted aryl, wherein the substituents include one or more alkyl or substituted alkyl groups or one or more nitro groups,

m is 0 or 1,

n , p , and q are independently 0, 1, or 2,

25

provided that at least one of m , n , p , and q is other than 0;

X is O, S or N-alk,

30

R_5 and R_6 are independently hydrogen, alkyl, substituted alkyl, or taken together form a lower alkenyl ring of 5 or 6 members,

provided that if R_2 , R_3 , and R_4 are hydrogen,
then, both R_5 and R_6 may be neither hydrogen nor methyl;
further provided that if R_2 and R_3 are hydrogen while R_4 is trifluoromethyl
or if R_2 and R_4 are hydrogen while R_3 is chloro, then both R_5 and R_6 may not
be methyl;

R_7 is hydrogen, alkyl, $N(alk)_2$, substituted alkyl or OH (and the resulting tautomeric
form in which the OH is tautomerized to a carbonyl and the imine is tautomerized
to an NH group);

Y is nitrogen, CH, C-CN or C-C(O)OR₈; and
wherein R_8 is hydrogen, alkyl or substituted alkyl.

2. The use of claim 1, wherein the cardioprotection involves short term protection
during surgical procedures selected from the group consisting of percutaneous angioplasty
(PTDA), angioplasty, and cardiac surgeries.

3. The use of claim 1, wherein the cardioprotection involves long term protection
from myocardial infarction.

4. The use of claim 1, wherein the neuroprotection involves stroke prevention,
stroke treatment, or the treatment of epilepsy.

5. The use of claim 1, wherein the pain management involves the treatment of
diabetic neuropathy, post herpetic neuralgia or other forms of neuropathic pain.

6. The use of claim 5 wherein the compound is present in a suitable carrier for
acute i.v. injection, chronic oral administration or chronic intravenous injection.

7. The use of claim 1, wherein the treatment of diabetes includes the treatment of insulin and non-insulin dependent diabetes mellitus, the stimulation of insulin secretion from the pancreas, or the increase in tissue sensitivity to insulin.

5 8. The use of claim 1, wherein the treatment of GI disorders involves treating a disorder selected from the group consisting of diarrhea, irritable bowel disease, irritable bowel syndrome, and incontinence.

10 9. The use of claim 1 wherein the compound is selected from compounds of formula (IA).

10. The use of claim 1 wherein the compound is selected from compounds of formula (IB).

15 11. The use of claim 1 wherein the compound is selected from compounds of formula (IC).

12. The use of claim 1 wherein:

20 R_1 is hydrogen,

R_2 , R_3 , and R_4 are independently hydrogen, halogen, or trifluoromethyl,

t is 0, 1, 2, or 3,

25 Z is NH, $N-(CH_2)_{1-3}$ phenyl, N -(ethoxycarbonylmethyl), N -(2- t -butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl), N -(3-methylbut-2-en-1-yl), N -(4-methylphenylsulphonyl), N -(4-nitro-(2-phenyleth-1-yl) or N -(benzyloxycarbonyl);

30

R₅ and R₆ are both hydrogen or both methyl, or R₅ and R₆ together form a cyclopentyl or cyclohexyl ring;

R₇ is hydrogen or methyl; and

5

R₈ is ethyl.


13. The use of claim 1 wherein the compound is selected from the group consisting of:

- 10 (2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-
phenyl]methanone,
(2-amino-3-thienyl)-(4-chlorophenyl)methanone,
2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
15 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(3-phenylprop-1-yl)-4,5,6,7-
20 tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(phenylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(phenylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
25 2-amino-3-(4-chloro-benzoyl)-6-(2-phenyleth-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(fluoromethyl)-benzoyl]-6-(2-phenyleth-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(3-phenylprop-1-yl)-4,5,6,7-
30 tetrahydrothieno[2,3-c]pyridine,


- 2-amino-3-benzoyl-6-methyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(ethoxycarbonylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-(ethoxycarbonylmethyl)-4,5,6,7-tetrahydrothieno
5 [2,3-c]pyridine,
2-amino-3-benzoyl-6-(3-methylbut-2-en-yl) -4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
10 2-amino-3-benzoyl-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-[2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-
propion-1-yl]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-4,5,6,7-tetrahydrobenzo[b]thiophene,
15 4-phenyl-5,6,7,8-tetrahydro[1]benzothieno[2,3-d]pyrimidine,
2-methyl,3-ethoxycarbonyl-4-phenyl-5,6,7,8-tetrahydro[1]Benzothieno
[2,3-b]pyridine,
2-amino-3-(4-bromobenzoyl)-cyclopenta[b]thiophene,
2-amino-3-benzoyl-6-(4-methylphenylsulphonyl)-4,5,6,7-tetrahydrothieno-
20 [2,3-c]pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
25 2-amino-3-(4-chlorobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-iodobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
30 2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,

- 2-amino-3-(4-bromobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
5 2-amino-3-(4-iodobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno(2,3-
c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6,6-bis(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
10 (2,3-c)pyridinium chloride,
2-amino-3-(4-fluorobenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6,6-bis(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno
15 (2,3-c)pyridinium chloride,
2-amino-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine, and
2-hydroxy-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine.


14. A method of determining whether a compound is active at moderating
20 adenosine receptors in a mammal, comprising performing competitive binding studies with
the compound to be analyzed and a compound selected from a group consisting of
compounds of formulas IA, IB, and IC:



(IA)



(IB)



(IC)

wherein:

R₁ is hydrogen, alkyl, substituted alkyl, or haloacetyl;

R₂, R₃, and R₄ are independently hydrogen, halogen, alkyl, substituted alkyl, aryl,
 5 lower alkenyl, lower alkanoyl, amino, trifluoromethyl, amino lower alkyl, nitro, or
 cyano;

t is 0, 1, 2, or 3;

Z is NH, N-(Gr)_m(Am)_n(Alk)_p(Ar)_q, or CH-(Gr)_m(Am)_n(Alk)_p(Ar)_q,

10

wherein

Gr is -SO₂-, -C(O)O-, or -C(O)-,

Am is -CH(NH₂)-, an amino acid residue, or an amino protected
 amino acid residue,

Alk is hydrogen, alkylene, substituted alkylene, alkenylene or substituted alkenylene,

Ar is aryl,

m is 0 or 1,

n, p, and q are independently 0, 1, or 2,

provided that at least one of m, n, p, and q is other than 0;

R₅ and R₆ are independently hydrogen, alkyl, substituted alkyl, or taken together form a lower alkenyl ring of 5 or 6 members,

provided that if R₂, R₃, and R₄ are hydrogen,

then, both R₅ and R₆ may be neither hydrogen nor methyl;

further provided that if R₂ and R₃ are hydrogen while R₄ is trifluoromethyl or if R₂ and R₄ are hydrogen while R₃ is chloro, then both R₅ and R₆ may not be methyl;

R₇ is hydrogen, alkyl or substituted alkyl;

Y is nitrogen, CH or C-C(O)OR₈; and

R₈ is hydrogen, alkyl or substituted alkyl.

15. The method of claim 14 wherein the compound is selected from compounds of formula (IA).

16. The method of claim 14 wherein the compound is selected from compounds of formula (IB).

17. The method of claim 14 wherein the compound is selected from compounds of formula (IC).

18. The method of claim 14 wherein:

5

R_1 is hydrogen,

R_2 , R_3 , and R_4 are independently hydrogen, halogen, or trifluoromethyl,

10

t is 0, 1, 2, or 3,

Z is NH, N-(CH₂)₁₋₃ phenyl, N-(ethoxycarbonylmethyl), N-(2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl), N-(3-methylbut-2-en-1-yl), N-(4-methylphenylsulphonyl), N-(4-nitro-(2-phenyleth-1-yl) or N-(benzyloxycarbonyl);

15

R_5 and R_6 are both hydrogen or both methyl, or R_5 and R_6 together form a cyclopentyl or cyclohexyl ring;

20

R_7 is hydrogen or methyl; and

R_8 is ethyl.

19. The method of claim 14 wherein the compound is selected from the group consisting of:

25

(2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-phenyl)]methanone,

(2-amino-3-thienyl)-(4-chlorophenyl)methanone,

2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno

30

[2,3-c]pyridine,

- 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
5 2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(3-phenylprop-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(phenylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(phenylmethyl)-4,5,6,7-
10 tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(2-phenyleth-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(fluoromethyl)-benzoyl]-6-(2-phenyleth-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
15 2-amino-3-(4-chloro-benzoyl)-6-(3-phenylprop-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-methyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(ethoxycarbonylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
20 2-amino-3-benzoyl-6-(ethoxycarbonylmethyl)-4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
2-amino-3-benzoyl-6-(3-methylbut-2-en-yl) -4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-
25 tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-[2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-
propion-1-yl]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
30 2-amino-3-benzoyl-4,5,6,7-tetrahydrobenzo[b]thiophene,

- 4-phenyl-5,6,7,8-tetrahydro[1]benzothieno[2,3-d]pyrimidine,
2-methyl,3-ethoxycarbonyl-4-phenyl-5,6,7,8-tetrahydro[1]Benzothieno
[2,3-b]pyridine,
2-amino-3-(4-bromobenzoyl)-cyclopenta[b]thiophene,
5 2-amino-3-benzoyl-6-(4-methylphenylsulphonyl)-4,5,6,7-tetrahydrothieno-
[2,3-c]pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
10 (2,3-c)pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-iodobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
15 (2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
20 2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-iodobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
25 2-amino-3-(4-phenylbenzoyl)-6,6-bis(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridinium chloride,
2-amino-3-(4-fluorobenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
30 2-amino-3-(4-phenylbenzoyl)-6,6-bis(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno

(2,3-c)pyridinium chloride,
2-amino-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine, and
2-hydroxy-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine.

20. A compound selected from the group consisting of:

- 5 (2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-
phenyl]]methanone,
(2-amino-3-thienyl)-(4-chlorophenyl)methanone,
2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
10 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(3-phenylprop-1-yl)-4,5,6,7-
15 tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(phenylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(phenylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
20 2-amino-3-(4-chloro-benzoyl)-6-(2-phenyleth-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-[3-(fluoromethyl)-benzoyl]-6-(2-phenyleth-1-yl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(3-phenylprop-1-yl)-4,5,6,7-
25 tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-methyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-(ethoxycarbonylmethyl)-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-(ethoxycarbonylmethyl)-4,5,6,7-tetrahydrothieno
30 [2,3-c]pyridine,

- 2-amino-3-benzoyl-6-(3-methylbut-2-en-yl)-4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
5 2-amino-3-benzoyl-6-[4-nitro-(2-phenyleth-1-yl)]-4,5,6,7-
tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-6-[2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-
propion-1-yl]-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-benzoyl-4,5,6,7-tetrahydrobenzo[b]thiophene,
10 4-phenyl-5,6,7,8-tetrahydro[1]benzothieno[2,3-d]pyrimidine,
2-methyl,3-ethoxycarbonyl-4-phenyl-5,6,7,8-tetrahydro[1]Benzothieno
[2,3-b]pyridine,
2-amino-3-(4-bromobenzoyl)-cyclopenta[b]thiophene,
2-amino-3-benzoyl-6-(4-methylphenylsulphonyl)-4,5,6,7-tetrahydrothieno-
15 [2,3-c]pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-chlorobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
20 2-amino-3-(4-chlorobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-iodobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(((phenyl)amino)carbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
25 2-amino-3-(4-bromobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-bromobenzoyl)-6-(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-iodobenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
30 2-amino-3-(4-iodobenzoyl)-6-(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno

- (2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6,6-bis(3-methyl-but-2-en-1-yl)-4,5,6,7-tetrahydrothieno
5 (2,3-c)pyridinium chloride,
2-amino-3-(4-fluorobenzoyl)-6-(benzyloxycarbonyl)-4,5,6,7-tetrahydrothieno
(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-4,5,6,7-tetrahydrothieno(2,3-c)pyridine,
2-amino-3-(4-phenylbenzoyl)-6,6-bis(prop-2-en-1-yl)-4,5,6,7-tetrahydrothieno
10 (2,3-c)pyridinium chloride,
2-amino-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine, and
2-hydroxy-3-cyano-4-phenyl-5,6,7,8-tetrahydro(1)benzothieno(2,3-b)pyridine.

21. A method for providing cardioprotection, neuroprotection, pain management,
reduction of free fatty acids, triglycerides, or glucose levels, adjunct therapy in diabetes,
15 treatment of GI disorders, treatment of glaucoma; treatment of sleep disorders; treatment of
cardiac disarrhythmias (peroxysmal supraventricular tachycardia, treatment of congestive
heart failure or treatment of inflammation, comprising administering to a patient in need of
treatment thereof an effective amount to treat the disorder of a compound as in claim 1.

1/16

CHO-HuA1 Membranes

Figure 1 (a)





Figure 1 (b)



Comp. 20 concentration (log M)

SUBSTITUTE SHEET (RULE 26)

CHO-HuA1 Membranes

Figure 2 (a)




Figure 2 (b)




Figure 2 (c)





Figure 2 (d)



CHO-HuA1 Membranes

Figure 3 (a)




Figure 3 (b)




Figure 3 (c)





Figure 3 (d)



4/16
Figure 4 (a)






Figure 4 (b)



5/16

Figure 5



6/16

Figure 6






Figure 7



8/16

Figure 8



9/16

Figure 9






Figure 10




11/16

Figure 11




12/16

Figure 12




13/16

Figure 13




14/16

Figure 14




15/16

Figure 15



16/16

Figure 16







INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/505, 31/435, 31/38, C07D 495/04, 333/78, 333/36, 333/66 // (C07D 495/04, 333:00, 239:00), (C07D 495/04, 333:00, 221:00)		A3	(11) International Publication Number: WO 99/21617
			(43) International Publication Date: 6 May 1999 (06.05.99)
(21) International Application Number: PCT/US98/20333		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 30 September 1998 (30.09.98)		Published With international search report.	
(30) Priority Data: 08/959,758 29 October 1997 (29.10.97) US 09/156,077 17 September 1998 (17.09.98) US		(88) Date of publication of the international search report: 29 July 1999 (29.07.99)	
(71) Applicant (for all designated States except US): MEDCO RESEARCH, INC. [US/US]; Suite 200, 85 T. W. Alexander Drive, Research Triangle Park, NC 27709 (US).			
(72) Inventor; and (75) Inventor/Applicant (for US only): BARALDI, Pier, Giovanni [IT/IT]; Via Tulipani, 73, I-66100 Ferrara (IT).			
(74) Agents: SAVAGE, Michael, G. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).			


(54) Title: ALLOSTERIC ADENOSINE RECEPTOR MODULATORS



(IA)



(IB)



(IC)

(57) Abstract.

The present invention relates to compounds of formulas (IA, IB and IC), the preparation thereof, pharmaceutical formulations thereof, and their use in medicine as allosteric adenosine receptor modulators for uses including protection against hypoxia and ischemia induced injury and treatment of adenosine-sensitive cardiac arrhythmias.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/20333

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/505 A61K31/435 A61K31/38 C07D495/04 C07D333/78
C07D333/36 C07D333/66
239:00), (C07D495/04, 333:00, 221:00)
/(C07D495/04, 333:00,

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BRUNS, R.F. ET AL.: "Structure-Activity Relationships for Enhancement of Adenosine A1 Receptor Binding by 2-Amino-3-Benzoylthiophenes" MOLECULAR PHARMACOLOGY, vol. 38, no. 6, 1990, pages 950-958, XP002093490 see abstract see Results see tables 1-3,6 see Discussion see page 958, column 1, last sentence --- -/--	1-10,12, 14

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

22 February 1999

Date of mailing of the international search report

28 05 99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Taylor, G

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/20333

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BRUNS R.F. ET AL.: "Allosteric Enhancement of Adenosine A1 Receptor Binding and Function by 2-Amino-3-Benzoylthiophenes" MOLECULAR PHARMACOLOGY, vol. 38, no. 6, 1990, pages 939-949, XP002093491 see abstract see Results see figure 1 see page 949, column 1, paragraph 3 to the end of the document	1-10,12, 14
X	--- MIZUMURA T. ET AL.: "PD 81,723, an Allosteric Enhancer of the A1 Adenosine Receptor, Lowers the Threshold for Ischemic Preconditioning in Dogs" CIRCULATION RESEARCH, vol. 79, no. 3, 1996, pages 415-423, XP002093492 see abstract see page 415, column 1 - column 2 see page 421, discussion see page 423, last paragraph	1-10,12, 14
X	--- CAO X. AND PHILLIS, J.W.: "Adenosine A1 Receptor Enhancer, PD 81,723, and Cerebral Ischemia/Reperfusion Injury in the Gerbil" GEN. PHARMAC., vol. 26, no. 7, 1995, pages 1545-1548, XP002093493 see abstract see Introduction see Results see Discussion see Summary -----	1-10,12, 14

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 20333

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-8 (in part), 9, 10, 12-21 (in part)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 98/20333

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-8 (in part), 9, 10, 12-21 (in part)




The use of compounds (IA) or (IB).

2. Claims: 1-8 (in part), 11, 12-21 (in part)

The use of compounds (IC).



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 6 : A61K 31/505, 31/435, 31/38, C07D 495/04, 333/78, 333/36, 333/66 // (C07D 495/04, 333:00, 239:00), (C07D 495/04, 333:00, 221:00)</p>	A3	<p>(11) International Publication Number: WO 99/21617</p> <p>(43) International Publication Date: 6 May 1999 (06.05.99)</p>
<p>(21) International Application Number: PCT/US98/20333</p> <p>(22) International Filing Date: 30 September 1998 (30.09.98)</p> <p>(30) Priority Data: 08/959,758 29 October 1997 (29.10.97) US 09/156,077 17 September 1998 (17.09.98) US</p> <p>(71) Applicant (for all designated States except US): MEDCO RESEARCH, INC. [US/US]; Suite 200, 85 T. W. Alexander Drive, Research Triangle Park, NC 27709 (US).</p> <p>(72) Inventor; and (75) Inventor/Applicant (for US only): BARALDI, Pier, Giovanni [IT/IT]; Via Tulipani, 73, I-66100 Ferrara (IT).</p> <p>(74) Agents: SAVAGE, Michael, G. et al.; Burns, Doane, Swecker & Mathis, L.L.P., P.O. Box 1404, Alexandria, VA 22313-1404 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report With amended claims.</p> <p>(88) Date of publication of the international search report: 29 July 1999 (29.07.99)</p> <p>Date of publication of the amended claims: 10 September 1999 (10.09.99)</p>
<p>(54) Title: ALLOSTERIC ADENOSINE RECEPTOR MODULATORS</p> <div style="display: flex; justify-content: space-around; align-items: flex-end;"> <div style="text-align: center;">  <p>(IA)</p> </div> <div style="text-align: center;">  <p>(IB)</p> </div> <div style="text-align: center;">  <p>(IC)</p> </div> </div> <p>(57) Abstract</p> <p>The present invention relates to compounds of formulas (IA, IB and IC), the preparation thereof, pharmaceutical formulations thereof, and their use in medicine as allosteric adenosine receptor modulators for uses including protection against hypoxia and ischemia induced injury and treatment of adenosine-sensitive cardiac arrhythmias.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

AMENDED CLAIMS

[received by the International Bureau on 27 July 1999 (27.07.99);
original claims 12 and 18 amended; remaining claims unchanged (3 pages)]

7. The use of claim 1, wherein the treatment of diabetes includes the treatment of insulin and non-insulin dependent diabetes mellitus, the stimulation of insulin secretion from the pancreas, or the increase in tissue sensitivity to insulin.

5

8. The use of claim 1, wherein the treatment of GI disorders involves treating a disorder selected from the group consisting of diarrhea, irritable bowel disease, irritable bowel syndrome, and incontinence.

10

9. The use of claim 1 wherein the compound is selected from compounds of formula (IA).

10. The use of claim 1 wherein the compound is selected from compounds of formula (IB).

15

11. The use of claim 1 wherein the compound is selected from compounds of formula (IC).

12. The use of claim 1 wherein:

20

R₁ is hydrogen,

R₂, R₃, and R₄ are independently hydrogen, halogen, or trifluoromethyl,

25

t is 0, 1, 2, or 3,

Z is NH, N-(ethoxycarbonylmethyl), N-(2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl), N-(3-methylbut-2-en-1-yl), N-(4-methylphenylsulphonyl), N-(4-nitro-(2-phenyleth-1-yl) or N-(benzyloxycarbonyl);

30

R₅ and R₆ are both hydrogen or both methyl;

R₇ is hydrogen or methyl; and

5 R₈ is ethyl.

13. The use of claim 1 wherein the compound is selected from the group consisting of:

- 10 (2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-phenyl]methanone,
- (2-amino-3-thienyl)-(4-chlorophenyl)methanone,
- 2-amino-3-benzoyl-6-benzoyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 15 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 2-amino-3-(4-chloro-benzoyl)-6-benzoyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 2-amino-3-(4-chloro-benzoyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(3-phenylprop-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 20 2-amino-3-(4-chloro-benzoyl)-6-(phenylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 2-amino-3-[3-(trifluoromethyl)-benzoyl]-6-(phenylmethyl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 2-amino-3-(4-chloro-benzoyl)-6-(2-phenyleth-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 25 2-amino-3-[3-(fluoromethyl)-benzoyl]-6-(2-phenyleth-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
- 2-amino-3-(4-chloro-benzoyl)-6-(3-phenylprop-1-yl)-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,

17. The method of claim 14 wherein the compound is selected from compounds of formula (IC).

18. The method of claim 14 wherein:

5 R_1 is hydrogen,

R_2 , R_3 , and R_4 are independently hydrogen, halogen, or trifluoromethyl,

10 t is 0, 1, 2, or 3,

Z is NH, N-(ethoxycarbonylmethyl), N-(2-t-butoxycarbonylamino-3-(4-hydroxyphenyl)-propion-1-yl), N-(3-methylbut-2-en-1-yl), N-(4-methylphenylsulphonyl), N-(4-nitro-(2-phenyleth-1-yl) or N-

15 (benzyloxycarbonyl);

R_5 and R_6 are both hydrogen or both methyl;

R_7 is hydrogen or methyl; and

20

R_8 is ethyl.

19. The method of claim 14 wherein the compound is selected from the group consisting of:

25 (2-amino-4,5-dimethyl-3-thienyl)-[(3,5-dichloro-4-amino)-phenyl]methanone,
(2-amino-3-thienyl)-(4-chlorophenyl)methanone,
2-amino-3-benzoyl-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno
[2,3-c]pyridine,
30 2-amino-3-benzoyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,
2-amino-3-(4-chloro-benzoyl)-6-benzyloxycarbonyl-4,5,6,7-tetrahydrothieno[2,3-c]pyridine,